

A

[This question paper contains 6 printed pages.]

Your Roll No.....

Sr. No. of Question Paper : 1382

C

Unique Paper Code : 32221301

Name of the Paper : Mathematical Physics – II

Name of the Course : B.Sc. (H) Physics

Semester : III

Duration : 3 Hours Maximum Marks : 75

**Instructions for Candidates**

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. Attempt **five** questions in all.
3. Q. No. 1 is compulsory.

1. Attempt any **five** questions :  $(5 \times 3 = 15)$

(a) Prove that even function can have no sine terms in its Fourier expansion.

P.T.O.

(b) Determine whether the functions  $\cos 2x$  and  $\cos x$  are orthogonal or not in the interval  $(0, 2\pi)$ .

(c) Evaluate :  $\int_0^{\pi/2} \cos^6 \theta d\theta$ .

(d) Find the value of  $\Gamma\left(\frac{-5}{2}\right)$ .

(e) Show that for integral values of  $n$ ,  $AJ_n(x) + BJ_{-n}(x)$  is not a general solution of Bessel equation of order  $n$ .

(f) Prove :  $P'_n(1) = \frac{n(n+1)}{2}$ .

(g) Find whether  $x = 1$  is an ordinary, regular or irregular singular point of the given differential equation :

$$x^2(1-x^2)y'' + \frac{2}{x}y' + 4y = 0$$

(h) Determine whether or not  $u(x, y) = 4e^{-3x} \cos 3y$  is a solution of given partial differential equation :

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

2. (a) Find the Fourier series expansion of a periodic function given by : (10)

$$f(t) = \begin{cases} E_0 \sin t, & 0 < t < \pi \\ 0, & \pi < t < 2\pi \end{cases}$$

(b) Evaluate :  $\int_0^2 x(8-x^3)^{1/3} dx$  (5)

3. Consider a periodic function  $f(x)$  of period  $2\pi$  such that

$$f(x) = \pi - x, \quad 0 < x < \pi$$

- (a) Plot odd extension of  $f(x)$  in the range  $(-3\pi, 3\pi)$ . (3)

- (b) Find its half-range Fourier Sine Series. (6)

(c) Show that  $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}$  (3)

(d) Also, prove that  $1 + \frac{1}{4} + \frac{1}{9} + \dots = \frac{\pi^2}{6}$  (3)

4. Consider the following differential equation :

$$4x^2 y'' + 4xy' + (x^2 - 1)y = 0$$

P.T.O.

(a) Find whether  $x = 0$  is an ordinary, regular or irregular singular point. (3)

(b) Using Frobenius method, determine the roots of indicial equation and hence find the first solution. (4,5)

(c) Also, find the second solution. (3)

5. (a) Prove that orthogonality relation for Legendre polynomials is given by

$$\int_{-1}^1 P_n(x) P_m(x) dx = \begin{cases} \frac{2}{2n+1}, & m = n \\ 0, & m \neq n \end{cases} \quad (10)$$

(b) The generating function of Legendre polynomials is given by :

$$(1 - 2xt + t^2)^{-1/2} = \sum_{n=0}^{\infty} t^n P_n(x),$$

Using this generating function, prove that :

$$P_{n+1}(x) = \frac{2n+1}{n+1} x P_n(x) - \frac{n}{n+1} P_{n-1}(x) \quad (5)$$

6. Given,  $e^{\frac{x}{2}(t-\frac{1}{t})} = \sum_{n=-\infty}^{\infty} t^n J_n(x)$

Verify that :

$$(i) \cos(x\sin\theta) = J_0(x) + 2J_2(x) \cos 2\theta + 2J_4(x) \cos 4\theta + \dots$$

$$(ii) \sin(x\sin\theta) = 2J_1(x)\sin\theta + 2J_3(x)\sin 3\theta + \dots$$

Hence prove that :

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\theta - x\sin\theta) d\theta \quad (3,3,9)$$

7. Using the method of separation of variables, find the general solution of 2-D wave equation for the case of symmetrical circular membrane (radius = a):

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2}; c > 0$$

subject to the conditions :

$$u(a, t) = 0, \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = 0 \quad \text{and} \quad u(r, 0) = u_0(r) \quad (15)$$

8. (a) Using the method of separation of variables, solve the following differential equation :

P.T.O.

$$\frac{\partial u}{\partial x} = 4 \frac{\partial u}{\partial y}$$

when  $u(0, y) = 8e^{-3y} + 4e^{-5y}$ . (5)

- (b) A homogeneous rod of conducting material of length 100 cm has its ends kept at zero temperature and the initial temperature is

$$u(x, 0) = \begin{cases} x, & 0 \leq x \leq 50 \\ 100 - x, & 50 \leq x \leq 100 \end{cases}$$

Using 1-D heat equation, find the temperature  $u(x, t)$  at any time. (10)